

## ACTIVATION OF STORE – OPERATED $\text{Ca}^{2+}$ ENTRY IN CISPLATIN RESISTANT LEUKEMIC CELLS AFTER TREATMENT WITH PHOTOEXCITED FULLERENE $\text{C}_{60}$ AND CISPLATIN

D. V. FRANSKEVYCH, I. I. GRYNKYUK, S. V. PRYLUTSKA, O. P. MATYSHEVSKA

Taras Shevchenko National University of Kyiv, Ukraine;  
e-mail: [dashaqq@gmail.com](mailto:dashaqq@gmail.com)

$\text{Ca}^{2+}$ -regulating system in cancer cells is suggested to be remodulated particularly by reduced store-operated  $\text{Ca}^{2+}$  entry (SOCE) through plasma membrane in order to maintain moderately reduced cytosolic  $\text{Ca}^{2+}$  concentration and to avoid apoptosis. The endoplasmic reticulum (ER)  $\text{Ca}^{2+}$  pool content and the size of SOCE in leukemic wild type (L1210) and resistant to cisplatin (L1210R) cells in control, after treatment with either cisplatin (1  $\mu\text{g}/\text{ml}$ ) or photoexcited fullerene  $\text{C}_{60}$  ( $10^{-5}$  M) alone, or their combination were estimated with the use of Indo-1 AM. The SOCE in resistant to cisplatin L1210R cells was found to be lower than in the wild-type cells. After treatment with cisplatin the decrease of thapsigargin (TG)-sensitive ER  $\text{Ca}^{2+}$  pool with no significant increase of SOCE was observed in L1210 cells, while no changes were detected in L1210R cells. Photoexcitation of intracellular accumulated fullerene  $\text{C}_{60}$  in the visible range of spectrum (410-700 nm) was accompanied by increase of SOCE not only in sensitive, but in resistant cells as well. In resistant L1210R cells treated with photoexcited  $\text{C}_{60}$  essential effect of cisplatin on  $\text{Ca}^{2+}$  homeostasis became obvious: the size of SOCE proved to be higher than after treatment with photoexcited  $\text{C}_{60}$  alone. The data obtained allow suggesting the influence of photoexcited  $\text{C}_{60}$  not only on  $\text{Ca}^{2+}$ -regulating system, but on those involved in controlling cisplatin entry into drug resistant cancer cells.

**Key words:** calcium, SOCE, leukemic cells, cisplatin, drug resistance, fullerene  $\text{C}_{60}$ .

$\text{Ca}^{2+}$  is a ubiquitous signaling messenger involved in cell cycle, differentiation, proliferation and apoptosis regulation. Eukaryotic cells regulate  $\text{Ca}^{2+}$  concentration in cytosol ( $[\text{Ca}^{2+}]_{\text{cyt}}$ ) by its release from intracellular stores (endoplasmic reticulum and mitochondria) or influx through plasma membrane  $\text{Ca}^{2+}$  channels. In non-excitabile cells the major mechanism of  $\text{Ca}^{2+}$  influx into intracellular space is store-operated calcium entry (SOCE). SOCE is controlled by filling the endoplasmic reticulum (ER)  $\text{Ca}^{2+}$  store. Its depletion stimulates the opening of plasma membrane store-operated  $\text{Ca}^{2+}$  channels (SOCCs), that allows refilling the ER  $\text{Ca}^{2+}$  pool. The canonical components of store-operated  $\text{Ca}^{2+}$  entry are the integral ER membrane protein STIM-1, which acts as a  $\text{Ca}^{2+}$  sensor and plasma membrane protein Orai1, which fulfils channel function after juxtaposition with STIM [1, 2].

The moderate increase of cytosolic  $\text{Ca}^{2+}$  level promotes proliferation, while high amplitude elevation of  $[\text{Ca}^{2+}]_{\text{cyt}}$  induced, in particular, by SOCE can be followed by uncontrolled increase of  $\text{Ca}^{2+}$  concentration in mitochondria and cell death by apoptosis [3, 4].

In cancer, normal relationships among extracellular, cytosolic, endoplasmic reticulum and mitochondrial  $\text{Ca}^{2+}$  become distorted and  $\text{Ca}^{2+}$ -dependent signaling pathways are deregulated in a way to promote cancer hallmarks such as enhanced proliferation, survival and invasion [5]. Cancer cells have remodulated  $\text{Ca}^{2+}$  buffering system which allows maintaining  $[\text{Ca}^{2+}]_{\text{cyt}}$  constantly at the relatively low level and to avoid apoptosis [6, 7]. Store-operated  $\text{Ca}^{2+}$  entry through plasma membrane and its release from ER in cancer cells are reduced due to increased expression of IP3R channels and decreased expres-

sion of ER Ca<sup>2+</sup> ATPase [8,9]. It is shown that prostate cancer cells, upon transition to more aggressive androgen-independent phenotype, downregulate their SOCE by decreasing the expression of the principal plasma membrane SOC-channel-forming subunit, ORAI1 protein [10],

That is why searching for agents, which could activate Ca<sup>2+</sup>-dependent apoptosis by restoring SOCE in cancer cells, is promising.

The widely used antitumor drug cisplatin (cis-Pt) could induce Ca<sup>2+</sup>-dependent apoptosis via IP3R-mediated increase of cytosolic Ca<sup>2+</sup> level, induction of ER stress and caspase-12 activation, but chemotherapeutic effect of cisplatin is limited by development of cancer cells multidrug resistance [11, 12]. Modulation of signaling pathways involved in development of drug resistance is considered to be a perspective approach for anticancer therapy optimization.

The representative of carbon nanostructures fullerene C<sub>60</sub> demonstrates unique physicochemical properties: nanosize, ability to penetrate into cytoplasm and compatibility with biological molecules [13-15]. Fullerene C<sub>60</sub> *per se* is biologically inert and becomes biologically active after its exposure to light. Due to extended  $\pi$ -conjugated system of molecular orbitals fullerene C<sub>60</sub> absorbs UV/visible light efficiently and is able to generate cytotoxic reactive oxygen species (ROS) [16, 17]. Taking into account that components of cells' Ca<sup>2+</sup> buffering system are sensitive to ROS [18, 19], redox regulation of their activity seems to be one of the ways to modulate Ca<sup>2+</sup> homeostasis in cancer cells and to activate apoptosis program.

As it was previously shown by us with the use of fluorescent labeled C<sub>60</sub> and confocal microscopy, leukemic cells could effectively uptake fullerene C<sub>60</sub> from the medium [20]. Combined treatment of leukemic cells with photoexcited fullerene C<sub>60</sub> and 1  $\mu$ g/ml cisplatin allowed us to enhance cytotoxic effect and to accelerate cell death of both sensitive and resistant to cisplatin cells as compared with each agent to be used alone [20, 21], but the biochemical mechanisms of this effect need further elucidation. A substantial increase of [Ca<sup>2+</sup>]<sub>cyt</sub> in leukemic cells which was detected 3 h after combined treatment with photoexcited C<sub>60</sub> and cisplatin [21] could be one of the determining factors in this effect.

The aim of this study was to estimate the values of ER Ca<sup>2+</sup> pool and store-operated calcium entry in sensitive and resistant to cisplatin leukemic cells in control and after treatment with either cisplatin or

photoexcited fullerene C<sub>60</sub> alone, or their combination.

## Materials and Methods

Two murine leukemic cell lines – wild type, sensitive (L1210) and resistant (L1210R) to cisplatin were obtained from the Bank of Cell Lines from Human and Animal Tissues of R. E. Kavetsky Institute of Experimental Pathology, Oncology and Radiobiology, NAS of Ukraine. The both cell lines were cultured in RPMI 1640 medium supplemented with 10% fetal bovine serum (FBS, Sigma-Aldrich, Germany), 50  $\mu$ g/ml penicillin and 100  $\mu$ g/ml streptomycin at 37 °C in a 5% CO<sub>2</sub> humidified atmosphere. Cisplatin in a range of 0.1-10  $\mu$ g/ml was shown to decrease the viability of L1210 in a dose dependent manner up to 90% during 72 h incubation with no effect on L1210R cells [21].

Homogeneous stable water colloid solution of fullerene C<sub>60</sub> (10<sup>-4</sup> M, purity >99.5%, nanoparticle average size 50 nm) was obtained at Technical University of Ilmenau (Germany) [22, 23]. Cells were incubated in RPMI 1640 medium for 2 h with or without fullerene C<sub>60</sub> (10<sup>-5</sup> M). Photoactivation of fullerene C<sub>60</sub> accumulated by cells was done by probes irradiation with light-emitting diode lamp (410-700 nm light, irradiance 100 mW, during 2 min) in 12-well plate. After irradiation cells were incubated for 3 h and used for estimation of [Ca<sup>2+</sup>]<sub>cyt</sub> and SOCE. In the case of combined treatment with photoexcited fullerene and cisplatin (Sigma, USA) the drug in concentration of 1  $\mu$ g/ml was added to incubation medium.

Free cytosolic Ca<sup>2+</sup> concentration was measured using fluorescent probe Indo-1 (Sigma, USA). Cells (3 $\times$ 10<sup>7</sup>/ml) in buffer A consisting of (mM): KCl – 5, NaCl – 120, CaCl<sub>2</sub> – 1, glucose – 10, MgCl<sub>2</sub> – 1, NaHCO<sub>3</sub> – 4, HEPES – 10, pH 7.4 were loaded with Indo-1AM (1 mM) in the presence of 0.05% Pluronic F-127 (Sigma, USA) for 40 min at 25 °C, then washed by centrifugation (600 g, 10 min) and resuspended in buffer A. Cells loaded with Indo-1 (2.5 $\times$ 10<sup>6</sup> cells/ml) were treated with C<sub>60</sub>, cisplatin or both as indicated above. Following incubation the cells were transferred to nominally Ca<sup>2+</sup>-free buffer A containing 0.1 mM EGTA. ER Ca<sup>2+</sup> content was estimated after addition of 1  $\mu$ M thapsigargin (TG), store-operated Ca<sup>2+</sup> entry was examined by adding 1 mM CaCl<sub>2</sub> to incubation medium. Indo-1 fluorescence in cells was recorded using spectrofluorometer (Shimadzu RF-510, Japan),  $\lambda$  excitation – 350 nm,

$\lambda$  emission – 410 and 495 nm. The concentration of free cytosolic  $\text{Ca}^{2+}$  was calculated as described in [24].

The data were represented as mean  $\pm$  SD of more than four independent experiments. Mean (M) and standard deviation (SD) were calculated for each group. Statistical analysis was performed using one-way ANOVA followed by post Tukey test. A value of  $P < 0.05$  was considered statistically significant. Data processing was performed by IBM PC using specialized applications GraphPad Prism 7 (GraphPad Software Inc., USA).

### Results and Discussion

The results of estimation of ER  $\text{Ca}^{2+}$  pool and store-operated calcium entry in sensitive and resistant to cisplatin L1210 cells in control are presented on Fig. 1 and Table. The ER  $\text{Ca}^{2+}$  pool content was estimated indirectly after Indo-1 loaded cells transfer into  $\text{Ca}^{2+}$ -free medium and addition of ER  $\text{Ca}^{2+}$ -ATPase inhibitor thapsigargin. The equilibrium concentration of cytosolic  $\text{Ca}^{2+}$  after 10 min incubation of L1210 and L1210R cells in  $\text{Ca}^{2+}$ -free medium was  $110 \pm 8$  and  $72 \pm 4$  nM respectively. It was shown that  $1 \mu\text{M}$  TG caused transient low amplitude increase of  $[\text{Ca}^{2+}]_{\text{cyt}}$ , indicating the depletion of ER calcium pool (Fig. 1). The TG-induced increase of  $[\text{Ca}^{2+}]_{\text{cyt}}$  above the equilibrium level at zero calcium ( $\Delta[\text{Ca}^{2+}]_{\text{cyt}}$ ) was considered as the measure of ER  $\text{Ca}^{2+}$  pool content (Table). Subsequent readmission of  $\text{Ca}^{2+}$  into incubation medium of TG-treated cells allowed examining extracellular  $\text{Ca}^{2+}$  influx pathway. The size of  $\text{Ca}^{2+}$ -induced increase of  $[\text{Ca}^{2+}]_{\text{cyt}}$  above




Fig. 1. Store-operated  $\text{Ca}^{2+}$  entry following thapsigargin-induced depletion of ER  $\text{Ca}^{2+}$  pool in sensitive and resistant to cisplatin L1210 cells

Table. ER  $\text{Ca}^{2+}$  pool and SOCE in leukemic cells

| Preparations   | L1210        | L1210R         |
|--|--------------|----------------|
| Thapsigargin-induced depletion of ER, $\Delta[\text{Ca}^{2+}]_{\text{cyt}}$ , nM | $190 \pm 7$  | $140 \pm 4^*$  |
| SOCE, $\Delta[\text{Ca}^{2+}]_{\text{cyt}}$ , nM                                 | $794 \pm 18$ | $595 \pm 20^*$ |

\* $P < 0.05$  in comparison with L1210 cells

the equilibrium level after TG addition was considered as the measure of SOCE. Addition of  $1 \text{ mM}$   $\text{CaCl}_2$  caused high amplitude increase of  $[\text{Ca}^{2+}]_{\text{cyt}}$ , which was smaller in L1210R cells than in L1210 cells (Fig. 1). The size of SOCE was found to be 1.3 times lower in resistant than in sensitive to cisplatin L1210 cells (Table).

The difference of SOCE value in resistant and sensitive to anticancer drug cells was also demonstrated in [25], where SOCE in resistant to cisplatin lung cancer cells A549 was found to be reduced as compared with wild-type cells A549. It is assumed that reduced activation of SOCE may contribute to the cisplatin resistant phenotype of L1210 cells [26].

As it was earlier shown by us, a substantial increase of  $[\text{Ca}^{2+}]_{\text{cyt}}$  in leukemic cells was detected 3 h after combined treatment with photoexcited  $\text{C}_{60}$  and cisplatin [21]. To investigate the possible reason of this event  $\text{Ca}^{2+}$  liberation from intracellular stores into the cytosol of leukemic cells was examined. The basal equilibrium concentration of cytosolic  $\text{Ca}^{2+}$  was estimated after cells transfer into  $\text{Ca}^{2+}$  free medium. It should be noted that fullerene  $\text{C}_{60}$  or light irradiation each applied separately had no evident effect on  $[\text{Ca}^{2+}]_{\text{cyt}}$ , TG-sensitive  $\text{Ca}^{2+}$  pool or SOCE (Figs. 2-4).

We have found that treatment of wild-type L1210 cells with cisplatin in concentration  $1 \mu\text{g/ml}$  was followed by the 1.7 times increase of basal equilibrium cytosolic  $\text{Ca}^{2+}$  concentration after incubation in  $\text{Ca}^{2+}$  free medium in comparison with control. No elevation of basal equilibrium  $[\text{Ca}^{2+}]_{\text{cyt}}$  in resistant to cisplatin L1210R cells was detected (Fig. 2).

In contrast to cisplatin, photoexcited fullerene  $\text{C}_{60}$  induced a considerable rise of basal equilibrium  $[\text{Ca}^{2+}]_{\text{cyt}}$  in cells of both lines (Fig. 2).

More pronounced increase of  $[\text{Ca}^{2+}]_{\text{cyt}}$  in the cells of both lines in the  $\text{Ca}^{2+}$  free medium was detected after combined action of photoexcited fullerene  $\text{C}_{60}$  and cisplatin. The level of cytosolic  $\text{Ca}^{2+}$  reached  $317 \pm 15$  and  $607 \pm 24$  nM in L1210 and L1210R cells, respectively (Fig. 2). The data obtained allowed us to suggest that enhancement of  $[\text{Ca}^{2+}]_{\text{cyt}}$




Fig. 2. Concentration of cytosolic  $\text{Ca}^{2+}$  (nM) in leukemic cells transferred into  $\text{Ca}^{2+}$ -free medium 3 h after treatment. \* $P < 0.05$  in comparison with control,  $\div P < 0.05$  in comparison with cisplatin, # $P < 0.05$  in comparison with photoexcited  $\text{C}_{60}$

in the  $\text{Ca}^{2+}$  free medium is the result of enhanced ER  $\text{Ca}^{2+}$  leak and/or reduced ER  $\text{Ca}^{2+}$  uptake. To confirm this the values of ER  $\text{Ca}^{2+}$  pool and store-operated calcium entry in leukemic cells 3 h after treatment with photoexcited fullerene  $\text{C}_{60}$ , cisplatin, and its combination were studied.

After treatment of wild-type L1210 cells with 1  $\mu\text{g}/\text{ml}$  cisplatin the TG-induced increase of  $[\text{Ca}^{2+}]_{\text{cyt}}$  above the equilibrium level was found to be somewhat reduced (Fig. 3) with no significant change of SOCE in comparison with control (Fig. 4). No change of either ER  $\text{Ca}^{2+}$ -content (Fig. 3) or SOCE (Fig. 4) was observed in L1210R after treatment with cisplatin.

After photoexcitation of accumulated fullerene  $\text{C}_{60}$  TG-sensitive ER  $\text{Ca}^{2+}$  pool was found to be decreased with simultaneous increase of SOCE both in sensitive and resistant to cisplatin leukemic cells. The TG-induced increase of  $[\text{Ca}^{2+}]_{\text{cyt}}$  above the equilibrium level in L1210 and L1210R was 3.5 and 2 times lower (Fig. 3), while the size of SOCE was 2.2 and 1.8 times higher than in control, respectively (Fig. 4). As it was previously shown, photoexcitation of fullerene  $\text{C}_{60}$  accumulated by human lymphoma Jurkat cells was also accompanied by enhancement of SOCE and cell death by apoptosis [27]. We assume that the observed enhancement of SOCE in leukemic cells of both lines could be mediated in part by intense ROS production after fullerene  $\text{C}_{60}$

photoexcitation [28]. Previous studies have revealed a cross-talk between ROS and components of  $\text{Ca}^{2+}$ -regulating system [29]. Thus, functional activity of both IP3R and STIM appeared to be dependent on oxidation of their reactive thiol groups. Oxidation of IP3R thiol groups was shown to be followed by activation of IP3 receptors and  $\text{Ca}^{2+}$  release from ER [30, 31], while  $\text{H}_2\text{O}_2$ -induced modification of STIM Cys56 residue was followed by SOCE intensification [32].

After combined treatment of L1210 cells with photoexcited fullerene  $\text{C}_{60}$  and cisplatin the additive effect of both agents was detected, the ER  $\text{Ca}^{2+}$  content, as probed by TG application, was lower (Fig. 3) and the size of SOCE was higher (Fig. 4) compared with the respective values after treatment with photoexcited  $\text{C}_{60}$  or cisplatin separately.

It is interesting that the evident effect of cisplatin on  $\text{Ca}^{2+}$  homeostasis characteristics in L1210R cells treated with photoexcited  $\text{C}_{60}$  became evident: the TG-sensitive ER  $\text{Ca}^{2+}$  pool was lower ( $48 \pm 4$  vs. of  $70 \pm 3$  nM) (Fig. 3) and the size of SOCE higher ( $1351 \pm 35$  vs.  $1042 \pm 40$ ) (Fig. 4). These data suggest that photoexcited  $\text{C}_{60}$  may act not only on components of  $\text{Ca}^{2+}$  regulating system, but also on those involved in controlling antitumor drug influx and accumulation in cancer cells.

Lower TG-sensitive ER  $\text{Ca}^{2+}$  pool content in the presence of photoexcited  $\text{C}_{60}$  and cisplatin could






Fig. 3. Thapsigargin sensitive ER  $Ca^{2+}$  pool in treated leukemic cells. \* $P < 0.05$  in comparison with control,  $\div P < 0.05$  in comparison with cisplatin, # $P < 0.05$  in comparison with photoexcited  $C_{60}$




Fig. 4. SOCE in treated leukemic cells. \* $P < 0.05$  in comparison with control,  $\div P < 0.05$  in comparison with cisplatin, # $P < 0.05$  in comparison with photoexcited  $C_{60}$

result in a higher values of SOCE and  $[Ca^{2+}]_{cyt}$  in leukemic cells. But it should be noted that the high level of  $[Ca^{2+}]_{cyt}$  in cells of both lines, particularly in L1210R cells transferred into  $Ca^{2+}$ -free medium after combined action of photoexcited  $C_{60}$  and cispla-

tin (Fig. 2) could be caused not only by leak of  $Ca^{2+}$  from the ER, but from mitochondria as well. This assumption is confirmed by the data of mitochondrial membrane potential dissipation in L1210 and L1210R cells after treatment with photoexcited  $C_{60}$  [20].

In summary, we could conclude that photo-excitation of accumulated fullerene  $C_{60}$  allowed to enhance cisplatin cytotoxic effects and activate  $Ca^{2+}$ -dependent apoptotic pathway not only in sensitive leukemic cells, but also in leukemic cells resistant to cisplatin.

### **АКТИВАЦІЯ ЄМНІСНОГО ВХОДУ $Ca^{2+}$ В РЕЗИСТЕНТНИХ ДО ЦИСПЛАТИНУ ЛЕЙКЕМІЧНИХ КЛІТИНАХ ЗА ДІЇ ФОТОЗБУДЖЕНОГО ФУЛЕРЕНУ $C_{60}$ ТА ЦИСПЛАТИНУ**

*Д. В. Франкевич, І. І. Гринюк,  
С. В. Прилуцька, О. П. Матишевська*

Київський національний університет  
імені Тараса Шевченка, Україна;  
e-mail: dashaqq@gmail.com

Припускають, що у злоякісно трансформованих клітинах  $Ca^{2+}$ -регуляторна система є ремодульованою, зокрема, внаслідок послаблення входу катіона крізь плазматичну мембрану за ємнісним механізмом (SOCE), що дозволяє утримувати концентрацію цитозольного  $Ca^{2+}$  на помірно підвищеному рівні та уникати апоптозу. У роботі з використанням зонда Indo-1AM оцінено відносні величини  $Ca^{2+}$ -пулу ендоплазматичного ретикулула та ємнісного входу  $Ca^{2+}$  у лейкемічних клітинах дикого типу (L1210) та резистентних до цисплатину (L1210R) у контролі, за дії цисплатину (1 мкг/мл) або фотозбудженого  $C_{60}$  ( $10^{-5}$  М) окремо, чи за їх комбінації. Показано, що величина SOCE в резистентних до цисплатину клітинах L1210R була нижчою, ніж у клітинах L1210. За дії цисплатину величина тапсигаргін (TG)-чутливого  $Ca^{2+}$  пулу ЕПР у клітинах L1210 знижувалась без підвищення SOCE, тоді як у клітинах L1210R не було відмічено змін. Фотозбудження акумульованого фулерену  $C_{60}$  у видимому діапазоні спектра (410–700 нм) спричиняло посилення SOCE не тільки в чутливих, але й в резистентних до дії цисплатину клітинах. У резистентних клітинах L1210R за дії фотозбудженого фулерену  $C_{60}$  цисплатин впливав на  $Ca^{2+}$  гомеостаз: величина SOCE зростала більше, ніж за дії тільки фотозбудженого  $C_{60}$ . Одержані дані вказують на вплив фотозбудженого  $C_{60}$  не тільки на  $Ca^{2+}$ -регуляторну систему, але й на компоненти систем, що контролюють надход-

ження цисплатину в резистентних до препарату злоякісно трансформованих клітинах.

**Ключові слова:** кальцій, SOCE, лейкемічні клітини, резистентність до цисплатину, фулерен  $C_{60}$ .

### **АКТИВАЦИЯ ЁМКОСТНОГО ВХОДА $Ca^{2+}$ В РЕЗИСТЕНТНЫХ К ЦИСПЛАТИНУ ЛЕЙКЕМИЧЕСКИХ КЛЕТКАХ ПРИ ДЕЙСТВИИ ФОТОВОЗБУЖДЕННОГО ФУЛЛЕРЕНА $C_{60}$ И ЦИСПЛАТИНА**

*Д. В. Франкевич, И. И. Гринюк,  
С. В. Прилуцкая, О. П. Матишевская*

Киевский национальный университет  
имени Тараса Шевченко, Украина;  
e-mail: dashaqq@gmail.com

Предполагается, что в злокачественно трансформированных клетках  $Ca^{2+}$ -регуляторная система ремодулирована, в частности, вследствие ослабленного входа катиона через плазматическую мембрану по ёмкостному механизму (SOCE), что позволяет поддерживать концентрацию цитозольного  $Ca^{2+}$  на умеренно повышенном уровне и избегать апоптоза. В работе с использованием зонда Indo-1AM оценены относительные величины  $Ca^{2+}$ -пула эндоплазматического ретикулула и ёмкостного входа  $Ca^{2+}$  в лейкемических клетках дикого типа (L1210) и резистентных к цисплатину (L1210R) в контроле, при действии цисплатина (1 мкг/мл) или фотозбудженного  $C_{60}$  ( $10^{-5}$  М) отдельно, или в комбинации. Показано, что величина SOCE в резистентных к цисплатину клетках L1210R была ниже, чем в клетках L1210. При действии цисплатина величина тапсигаргін (TG)-чувствительного  $Ca^{2+}$ -пула ЭПР в клетках L1210 снижалась без усиления SOCE, тогда как в клетках L1210R не было отмечено изменений. Фотозбуджение аккумуляированного фуллерена  $C_{60}$  в видимом диапазоне спектра (410–700 нм) приводило к усилению SOCE не только в чувствительных, но и в резистентных к цисплатину клетках. При действии фотозбудженного фуллерена  $C_{60}$  в резистентных L1210R клетках наблюдалось влияние цисплатина на  $Ca^{2+}$  гомеостаз: величина SOCE возросла больше, чем при действии только фотозбудженного  $C_{60}$ . Полученные данные указывают на

действие фотовозбужденного  $C_{60}$  не только на  $Ca^{2+}$ -регуляторную систему, но и на компоненты систем, контролирующих поступление цисплатина в резистентных к препарату злокачественно трансформированных клетках.

**Ключевые слова:** кальций, SOCE, лейкемические клетки, резистентность к цисплатину, фуллерен  $C_{60}$ .

### References

1. Parekh AB. Store-operated CRAC channels: function in health and disease. *Nat Rev Drug Discov.* 2010; 9(5): 399-410.
2. Park CY, Hoover PJ, Mullins FM, Bachhawat P, Covington ED, Raunser S, Walz T, Garcia KC, Dolmetsch RE, Lewis RS. STIM1 clusters and activates CRAC channels via direct binding of a cytosolic domain to Orai1. *Cell.* 2009; 136(5): 876-890.
3. Harr MW, Distelhorst CW. Apoptosis and autophagy: decoding calcium signals that mediate life or death. *Cold Spring Harb Perspect Biol.* 2010; 2(10): a005579.
4. Lebedzinska M, Szabadkai G, Jones AW, Duszynski J, Wieckowski MR. Interactions between the endoplasmic reticulum, mitochondria, plasma membrane and other subcellular organelles. *Int J Biochem Cell Biol.* 2009; 41(10): 1805-1816.
5. Prevarskaya N, Ouadid-Ahidouch H, Skryma R, Shuba Y. Remodelling of  $Ca^{2+}$  transport in cancer: how it contributes to cancer hallmarks? *Philos Trans R Soc Lond B Biol Sci.* 2014; 369(1638): 20130097.
6. Roderick HL, Cook SJ.  $Ca^{2+}$  signalling checkpoints in cancer: remodelling  $Ca^{2+}$  for cancer cell proliferation and survival. *Nat Rev Cancer.* 2008; 8(5): 361-375.
7. Yang H, Zhang Q, He J, Lu W. Regulation of calcium signaling in lung cancer. *J Thorac Dis.* 2010; 2(1): 52-56.
8. Kato H, Nishitoh H. Stress responses from the endoplasmic reticulum in cancer. *Front Oncol.* 2015; 5: 93.
9. Gélébart P, Kovács T, Brouland JP, van Gorp R, Grossmann J, Rivard N, Panis Y, Martin V, Bredoux R, Enouf J, Papp B. Expression of endomembrane calcium pumps in colon and gastric cancer cells. Induction of SERCA3 expression during differentiation. *J Biol Chem.* 2002; 277(29): 26310-26320.
10. Flourakis M, Lehen'kyi V, Beck B, Raphaël M, Vandenberghe M, Abeele FV, Roudbaraki M, Lepage G, Mauroy B, Romanin C, Shuba Y, Skryma R, Prevarskaya N. Orai1 contributes to the establishment of an apoptosis-resistant phenotype in prostate cancer cells. *Cell Death Dis.* 2010; 1: e75.
11. Galluzzi L, Vitale I, Michels J, Brenner C, Szabadkai G, Harel-Bellan A, Castedo M, Kroemer G. Systems biology of cisplatin resistance: past, present and future. *Cell Death Dis.* 2014; 5: e1257.
12. Köberle B, Tomicic MT, Usanova S, Kaina B. Cisplatin resistance: preclinical findings and clinical implications. *Biochim Biophys Acta.* 2010; 1806(2): 172-182.
13. Wilson S. R. Biological aspects of fullerenes. Fullerenes: Chemistry, Physics and Technology. New York: John Wiley & Sons. 2000; 437-465.
14. Piotrovsky LB, Kiselev OI. Fullerene in biology. SPb: Rostoc, 2006. 336 p. (In Russian).
15. Horie M, Nishio K, Kato H, Shinohara N, Nakamura A, Fujita K, Kinugasa S, Endoh S, Yamamoto K, Yamamoto O, Niki E, Yoshida Y, Iwahashi H. *In vitro* evaluation of cellular responses induced by stable fullerene  $C_{60}$  medium dispersion. *J Biochem.* 2010; 148(3): 289-298.
16. Yamakoshi Y, Umezawa N, Ryu A, Arakane K, Miyata N, Goda Y, Masumizu T, Nagano T. Active oxygen species generated from photoexcited fullerene ( $C_{60}$ ) as potential medicines:  $O_2^{*-}$  versus  $1O_2$ . *J Am Chem Soc.* 2003; 125(42): 12803-12809.
17. Moor KJ, Snow SD, Kim JH. Differential photoactivity of aqueous [C60] and [C70] fullerene aggregates. *Environ Sci Technol.* 2015; 49(10): 5990-5998.
18. Yan Y, Wei CL, Zhang WR, Cheng HP, Liu J. Cross-talk between calcium and reactive oxygen species signaling. *Acta Pharmacol Sin.* 2006; 27(7): 821-826.
19. Brookes PS, Yoon Y, Robotham JL, Anders MW, Sheu SS. Calcium, ATP, and ROS: a mitochondrial love-hate triangle. *Am J Physiol Cell Physiol.* 2004; 287(4): C817-C833.
20. Franskevych D, Palyvoda K, Petukhov D, Prylutska S, Grynyuk I, Schuetze C, Drobot L, Matyshevskaya O, Ritter U. Fullerene  $C_{60}$  Penetration into Leukemic Cells and Its Photoinduced Cytotoxic Effects. *Nanoscale Res Lett.* 2017; 12(1): 40.

21. Franskevych DV, Prylutska SV, Grynyuk II, Grebinyk DM, Matyshevska OP. Enhanced cytotoxicity of photoexcited fullerene C<sub>60</sub> and cisplatin combination against drug-resistant leukemic cells. *Exp Oncol.* 2015; 37(3): 187-191.
22. Scharff P, Risch K, Carta-Abelmann L, DmytrukIM, Bilyi MM, Golub OA, Khavryuchenko AV, Buzaneva EV, Aksenov VL, Avdeev MV, Prylutsky YuI, Durov SS. Structure of C<sub>60</sub> fullerene in water: spectroscopic data. *Carbon.* 2004; 42(5–6): 1203-1206.
23. Grynyuk II, Prylutska SV, Slobodyanik NS, Chunikhin OYu, Matyshevska OP. The aggregate state of C<sub>60</sub>-fullerene in various media. *Biotechnologia Acta.* 2013; 6(6): 71-76 (In Russian).
24. Gryniewicz G, Poenie M, Tsien RY. A new generation of Ca<sup>2+</sup> indicators with greatly improved fluorescence properties. *J Biol Chem.* 1985; 260(6): 3440-3450.
25. Jones A. Altered Cellular Signaling and Metabolism in Cisplatin Cytotoxicity and Chemoresistance. Available at <http://discovery.ucl.ac.uk/1335835/1/1335835.pdf>.
26. Hoffmann EK, Lambert IH. Ion channels and transporters in the development of drug resistance in cancer cells. *Philos Trans R Soc Lond B Biol Sci.* 2014; 369(1638): 20130109.
27. Grebinyk SM, Palyvoda KO, Prylutska SV, Grynyuk II, Samoylenko AA, Drobot LB, Matyshevska OP. Photoactivated fullerene C<sub>60</sub> induces store-operated Ca<sup>2+</sup> entry and cytochrome c release in Jurkat cells. *Ukr Biokhim Zhurn.* 2012; 84(6): 58-63.
28. Grynyuk I, Grebinyk S, Prylutska S, Mykhailova A, Franskevich D, Matyshevska O, Schutze C, Ritter U. Photoexcited fullerene C<sub>60</sub> disturbs prooxidant-antioxidant balance in leukemic L1210 cells. *Mat.-wiss. u. Werkstofftech.* 2013; 44(2–3): 139-143.
29. Csordás G, Hajnóczky G. SR/ER-mitochondrial local communication: calcium and ROS. *Biochim Biophys Acta.* 2009; 1787(11): 1352-1362.
30. Joseph SK, Nakao SK, Sukumvanich S. Reactivity of free thiol groups in type-I inositol trisphosphate receptors. *Biochem J.* 2006; 393(Pt 2): 575-582.
31. Foskett JK, White C, Cheung KH, Mak DO. Inositol trisphosphate receptor Ca<sup>2+</sup> release channels. *Physiol Rev.* 2007; 87(2): 593-658.
32. Bogeski I, Kilch T, Niemeyer BA. ROS and SOCE: recent advances and controversies in the regulation of STIM and Orai. *J Physiol.* 2012; 590(17): 4193-4200.

Received 10.10.2017